



# Answers

**E** Click here for exercises.

- $6t^5 + 4t^3 + 4t$
- $\frac{(1-t)(1-\sqrt{t})^2}{\sqrt{t}} - \frac{3(1-t)^2}{2\sqrt{t}}$
- $\frac{1}{\sqrt{1+t} + 1 + \sqrt{t}} \left( \frac{1}{2\sqrt{1+t}} + \frac{1+\sqrt{t}}{\sqrt{t}} \right)$
- $-e^{\cos t/e^{2t}} \left[ \left( 1 + \frac{\cos t}{e^{2t}} \right) \sin t - \frac{2e^{2t} \cos^2 t}{e^{4t}} \right]$
- $(18e^{2t} - 3 \cos t) e^t + (3e^t - 4 \cos t) \sin t$
- $e^{2t} \sqrt{1+e^{-2t}} (1+2t) - t \sqrt{1+e^{-2t}}$
- $y^2 z^3 (\cos t) + 2xyz^3 (-\sin t) + 3xy^2 z^2 (2e^{2t})$
- $\frac{1}{2y\sqrt{t}} + 2(\sin 2t) \left( \frac{x}{y^2} - \frac{1}{z} \right) + \frac{3y}{z^2 e^{3t}}$
- $4sx \sin y + 2tx^2 \cos y, 4xt \sin y + 2sx^2 \cos y$
- $2(s-t) \cos x \cos y - 2s \sin x \sin y,$   
 $2(t-s) \cos x \cos y + 2t \sin x \sin y$
- $(2x - 6xy^3) e^t - 9x^2 y^2 e^{-t}, (2x - 6xy^3) se^t + 9x^2 y^2 se^{-t}$
- $\frac{x^2 e^t}{1+x^2 y^2}, \left[ \tan^{-1}(xy) + \frac{xy}{1+x^2 y^2} \right] (2t) + \frac{x^2}{1+x^2 y^2} se^t$
- $(2^{x-3y} \ln 2) (2st - 3t^2), (2^{x-3y} \ln 2) (s^2 - 6st)$
- $(xe^y + e^{-x}) t^2, (e^y - ye^{-x}) e^t + 2(xe^y + e^{-x}) st$
- 2, 0
- 3, 2
- 0, 0, 4
- 0,  $-\frac{1}{4}, \frac{1}{2}$
- $-t/(p^2), 0, 1/p$
- $\sec(xy) [w + vzy \tan(xy)], z \sec(xy) \tan(xy) [yu + xw],$   
 $\sec(xy) [u + vzx \tan(xy)]$

**S** Click here for solutions.

- $st \sin(x-y) [2r \cos \theta - st^2 \sin \theta],$   
 $[rt \sin(x-y)] (r \cos \theta - 2st^2 \sin \theta),$   
 $[sr \sin(x-y)] (r \cos \theta - 3st^2 \sin \theta),$   
 $[-rst \sin(x-y)] (st^2 \cos \theta + r \sin \theta)$
- $2x - \frac{8x^2(x+2y)(x+y)}{y^{13/2}},$   
 $-8y + \frac{(x+2y)x^3 y^{1/2}(5x+2y)}{y^8}$
- $\frac{y-2x}{3y^2-x}$
- $-\frac{6xy^2+20x^3}{5y^4+6x^2y}$
- $\frac{y \sin x - \cos y}{\cos x - x \sin y}$
- $\frac{18x - x^{-2/3} y^{1/3}}{12y + x^{1/3} y^{-2/3}}$
- $\frac{z-y}{y-x}, \frac{x+z}{x-y}$
- $\frac{x-y-z}{z+x}, \frac{y-x}{z+x}$
- $-\frac{y^2 z^3 + 3x^2 y^2 z - 1}{3xy^2 z^2 + x^3 y^2 - 1}, -\frac{2xyz^3 + 2x^3 yz - 1}{3xy^2 z^2 + x^3 y^2 - 1}$
- $\frac{z \cos(xyz) - yze^{x+y}}{ye^{x+y} - x \cos(xyz)}, \frac{xz \cos(xyz) - e^{x+y}(2yz + y^2 z)}{y^2 e^{x+y} - xy \cos(xyz)}$
- $-\frac{y^2 + 2zx}{2yz + x^2}, -\frac{2xy + z^2}{2yz + x^2}$
- $-\frac{e^y + ze^x}{y + e^x}, -\frac{xe^y + z}{y + e^x}$
- $\frac{y^2 z^3 (x+yz) - 1}{y - 3xy^2 z^2 (x+yz)}, \frac{2xyz^3 (x+yz) - z}{y - 3xy^2 z^2 (x+yz)}$
- $-9600\pi \text{ cm}^3/\text{s}$

## Solutions

**E** Click here for exercises.

- $z = x^2 + y^2, x = t^3, y = 1 + t^2 \Rightarrow$   

$$\frac{dz}{dt} = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$$

$$= (2t^3)(3t^2) + 2(1 + t^2)(2t) = 6t^5 + 4t^3 + 4t$$
- $z = x^2 y^3, x = 1 + \sqrt{t}, y = 1 - \sqrt{t} \Rightarrow$   

$$\frac{dz}{dt} = 2xy^3 \frac{dx}{dt} + 3x^2 y^2 \frac{dy}{dt}$$

$$= 2xy^3 \frac{1}{2\sqrt{t}} + 3x^2 y^2 \left(-\frac{1}{2\sqrt{t}}\right)$$

$$= 2(1 + \sqrt{t})(1 - \sqrt{t})^3 \frac{1}{2\sqrt{t}}$$

$$+ 3(1 + \sqrt{t})^2(1 - \sqrt{t})^2 \left(-\frac{1}{2\sqrt{t}}\right)$$

$$= \frac{(1-t)(1-\sqrt{t})^2}{\sqrt{t}} - \frac{3(1-t)^2}{2\sqrt{t}}$$
- $z = \ln(x + y^2), x = \sqrt{1+t}, y = 1 + \sqrt{t} \Rightarrow$   

$$\frac{dz}{dt} = \frac{1}{(x + y^2)} \frac{1}{2\sqrt{1+t}} + \frac{1}{(x + y^2)} 2y \frac{1}{2\sqrt{t}}$$

$$= \frac{1}{\sqrt{1+t} + 1 + \sqrt{t}} \left( \frac{1}{2\sqrt{1+t}} + \frac{1 + \sqrt{t}}{\sqrt{t}} \right)$$
- $z = xe^{x/y}, x = \cos t, y = e^{2t} \Rightarrow$   

$$\frac{dz}{dt} = e^{x/y} \left( 1 + \frac{x}{y} \right) (-\sin t) - x^2 y^{-2} e^{x/y} (2e^{2t})$$

$$= -e^{\cos t / e^{2t}} \left[ \left( 1 + \frac{\cos t}{e^{2t}} \right) \sin t - \frac{2e^{2t} \cos^2 t}{e^{4t}} \right]$$
- $z = 6x^3 - 3xy + 2y^2, x = e^t, y = \cos t \Rightarrow$   

$$\frac{dz}{dt} = (18x^2 - 3y) e^t + (-3x + 4y) (-\sin t)$$

$$= (18e^{2t} - 3 \cos t) e^t + (3e^t - 4 \cos t) \sin t$$
- $z = x\sqrt{1-y^2}, x = te^{2t}, y = e^{-t} \Rightarrow$   

$$\frac{\partial z}{\partial t} = \sqrt{1+y^2} (e^{2t} + 2te^{2t})$$

$$+ \frac{1}{2}x(1+y^2)^{-1/2} (2y) (-e^{-t})$$

$$= e^{2t} \sqrt{1+e^{-2t}} (1+2t) - t\sqrt{1+e^{-2t}}$$
- $w = xy^2 z^3, x = \sin t, y = \cos t, z = 1 + e^{2t} \Rightarrow$   

$$\frac{dw}{dt} = y^2 z^3 (\cos t) + 2xy^2 z^3 (-\sin t) + 3xy^2 z^2 (2e^{2t})$$
- $w = \frac{x}{y} + \frac{y}{z}, x = \sqrt{t}, y = \cos 2t, z = e^{-3t} \Rightarrow$   

$$\frac{dw}{dt} = \frac{1}{y} \frac{1}{2\sqrt{t}} + \left( \frac{-x}{y^2} + \frac{1}{z} \right) (-2 \sin 2t) + \frac{-y}{z^2} (-3e^{-3t})$$

$$= \frac{1}{2y\sqrt{t}} + 2(\sin 2t) \left( \frac{x}{y^2} - \frac{1}{z} \right) + \frac{3y}{z^2 e^{3t}}$$

**A** Click here for answers.

- $z = x^2 \sin y, x = s^2 + t^2, y = 2st \Rightarrow$   

$$\frac{\partial z}{\partial s} = (2x \sin y) (2s) + (x^2 \cos y) (2t)$$

$$= 4sx \sin y + 2tx^2 \cos y$$

$$\frac{\partial z}{\partial t} = (2x \sin y) (2t) + (x^2 \cos y) (2s)$$

$$= 4xt \sin y + 2sx^2 \cos y$$
- $z = \sin x \cos y, x = (s-t)^2, y = s^2 - t^2 \Rightarrow$   

$$\frac{\partial z}{\partial s} = (\cos x \cos y) 2(s-t) - (\sin x \sin y) (2s)$$

$$= 2(s-t) \cos x \cos y - (2s) \sin x \sin y$$

$$\frac{\partial z}{\partial t} = (\cos x \cos y) (-2)(s-t) - (\sin x \sin y) (-2t)$$

$$= 2(t-s) \cos x \cos y + 2t \sin x \sin y$$
- $z = x^2 - 3x^2 y^3, x = se^t, y = se^{-t} \Rightarrow$   

$$\partial z / \partial s = (2x - 6xy^3) (e^t) + (-9x^2 y^2) (e^{-t})$$

$$= (2x - 6xy^3) e^t - 9x^2 y^2 e^{-t}$$

$$\partial z / \partial t = (2x - 6xy^3) (se^t) + (-9x^2 y^2) (-se^{-t})$$

$$= (2x - 6xy^3) se^t + 9x^2 y^2 se^{-t}$$
- $z = x \tan^{-1}(xy), x = t^2, y = se^t \Rightarrow$   

$$\frac{\partial z}{\partial s} = \left[ \tan^{-1}(xy) + \frac{x}{1+x^2 y^2} \right] (0) + \frac{x^2}{1+x^2 y^2} e^t$$

$$= \frac{x^2 e^t}{1+x^2 y^2}$$

$$\frac{\partial z}{\partial t} = \left[ \tan^{-1}(xy) + \frac{xy}{1+x^2 y^2} \right] (2t) + \frac{x^2}{1+x^2 y^2} se^t$$
- $z = 2^{x-3y}, x = s^2 t, y = st^2 \Rightarrow$   

$$\partial z / \partial s = (z \ln 2) (2st) + z (-3 \ln 2) (t^2)$$

$$= (2^{x-3y} \ln 2) (2st - 3t^2)$$

$$\partial z / \partial t = (z \ln 2) (s^2) + z (-3 \ln 2) (2st)$$

$$= (2^{x-3y} \ln 2) (s^2 - 6st)$$
- $z = xe^y + ye^{-x}, x = e^t, y = st^2 \Rightarrow$   

$$\partial z / \partial s = (e^y - ye^{-x}) (0) + (xe^y + e^{-x}) (t^2)$$

$$= (xe^y + e^{-x}) t^2$$

$$\partial z / \partial t = (e^y - ye^{-x}) (e^t) + (xe^y + e^{-x}) (2st)$$
- $w = x^2 + y^2 + z^2, x = st, y = s \cos t, z = s \sin t \Rightarrow$   

$$\frac{\partial w}{\partial s} = \frac{\partial w}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial s} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial s}$$

$$= 2xt + 2y \cos t + 2z \sin t$$

When  $s = 1, t = 0$ , we have  $x = 0, y = 1$  and  $z = 0$ , so  
 $\partial w / \partial s = 2 \cos 0 = 2$ . Similarly  
 $\partial w / \partial t = 2xs + 2y(-s \sin t) + 2z(s \cos t)$   
 $= 0 + (-2) \sin 0 + 0 = 0$   
when  $s = 1$  and  $t = 0$ .

16.  $u = xy + yz + zx, x = st, y = e^{st}, z = t^2 \Rightarrow$   
 $\partial u / \partial s = (y + z)t + (x + z)te^{st} + (x + y)(0)$  and  
 $\partial u / \partial t = (y + z)s + (x + z)se^{st} + (x + y)(2t)$ . When  
 $s = 0, t = 1$ , we have  $x = 0, y = 1, z = 1$ , so  
 $\partial u / \partial s = 2 + 1 + 0 = 3$  and  $\partial u / \partial t = 0 + 0 + (1)(2) = 2$ .

17.  $z = y^2 \tan x, x = t^2 uv, y = u + tv^2 \Rightarrow$   
 $\partial z / \partial t = (y^2 \sec^2 x) 2tuv + (2y \tan x) v^2$ ,  
 $\partial z / \partial u = (y^2 \sec^2 x) t^2 v + 2y \tan x$ ,  
 $\partial z / \partial v = (y^2 \sec^2 x) t^2 u + (2y \tan x) 2tv$ . When  $t = 2$ ,  
 $u = 1$  and  $v = 0$ , we have  $x = 0, y = 1$ , so  $\partial z / \partial t = 0$ ,  
 $\partial z / \partial u = 0, \partial z / \partial v = 4$ .

18.  $z = \frac{x}{y}, x = re^{st}, y = rse^t \Rightarrow$   
 $\frac{\partial z}{\partial r} = \frac{1}{y}e^{st} + \frac{-x}{y^2}se^t, \frac{\partial z}{\partial s} = \frac{1}{y}rte^{st} - \frac{x}{y^2}re^t$ ,  
 $\frac{\partial z}{\partial t} = \frac{1}{y}rse^{st} - \frac{x}{y^2}rse^t$ . When  $r = 1, s = 2$  and  $t = 0$ , we  
have  $x = 1, y = 2$ , so  $\partial z / \partial r = \frac{1}{2} + \frac{-1}{4} \cdot 2 = 0$ ,  
 $\partial z / \partial s = 0 - \frac{1}{4} = -\frac{1}{4}$  and  $\partial z / \partial t = \frac{1}{2} \cdot 2 - \frac{1}{4} \cdot 2 = \frac{1}{2}$ .

19.  $u = \frac{x+y}{y+z}, x = p+r+t, y = p-r+t, z = p+r-t \Rightarrow$   
 $\frac{\partial u}{\partial p} = \frac{1}{y+z} + \frac{(y+z) - (x+y)}{(y+z)^2} - \frac{x+y}{(y+z)^2}$   
 $= \frac{(y+z) + (z-x) - (x+y)}{(y+z)^2} = 2 \frac{z-x}{(y+z)^2}$   
 $= 2 \frac{-2t}{4p^2} = -\frac{t}{p^2}$   
 $\frac{\partial u}{\partial r} = \frac{1}{y+z} + \frac{z-x}{(y+z)^2}(-1) - \frac{x+y}{(y+z)^2} = 0$ , and  
 $\frac{\partial u}{\partial t} = \frac{1}{y+z} + \frac{z-x}{(y+z)^2} + \frac{x+y}{(y+z)^2}$   
 $= 2 \frac{y+z}{(y+z)^2} = \frac{2}{2p} = \frac{1}{p}$

20.  $t = z \sec(xy), x = uv, y = vw, z = wu \Rightarrow$   
 $\frac{\partial t}{\partial u} = [zy \sec(xy) \tan(xy)]v + [zx \sec(xy) \tan(xy)](0)$   
 $+ [\sec(xy)]w$   
 $= \sec(xy)[w + vzy \tan(xy)]$   
 $\frac{\partial t}{\partial v} = [zy \sec(xy) \tan(xy)]u + [zx \sec(xy) \tan(xy)]w$   
 $+ [\sec(xy)](0)$   
 $= z \sec(xy) \tan(xy)[yu + xw]$   
 $\frac{\partial t}{\partial w} = [zy \sec(xy) \tan(xy)](0) + [zx \sec(xy) \tan(xy)]v$   
 $+ [\sec(xy)]u$   
 $= \sec(xy)[u + vzx \tan(xy)]$

21.  $\frac{\partial w}{\partial r} = [-\sin(x-y)]s^2t^3 \sin \theta + [\sin(x-y)]2rst \cos \theta$   
 $= st \sin(x-y)[2r \cos \theta - st^2 \sin \theta]$   
 $\frac{\partial w}{\partial s} = [-\sin(x-y)]2rst^3 \sin \theta + [\sin(x-y)]r^2t \cos \theta$   
 $= [rt \sin(x-y)](r \cos \theta - 2st^2 \sin \theta)$   
 $\frac{\partial w}{\partial t} = [-\sin(x-y)]3rs^2t^2 \sin \theta + [\sin(x-y)]r^2s \cos \theta$   
 $= [sr \sin(x-y)](r \cos \theta - 3st^2 \sin \theta)$   
 $\frac{\partial w}{\partial \theta} = [-\sin(x-y)]rs^2t^3 \cos \theta + [\sin(x-y)](-r^2st \sin \theta)$   
 $= [-rst \sin(x-y)](st^2 \cos \theta + r \sin \theta)$

22.  $\frac{\partial u}{\partial x} = (q - 2pr^2s) + p + (-2p^2rs) \frac{1}{y^4}$   
 $+ (-p^2r^2)(2y^{3/2})$   
 $= p + q - 2pr^2s - \frac{2p^2rs}{y^4} - 2p^2r^2y^{3/2}$   
 $= 2x - \frac{8x^2(x+2y)(x+y)}{y^{13/2}}$   
 $\frac{\partial u}{\partial y} = (q - 2pr^2s)(2) + p(-2)$   
 $+ (-2p^2rs)\left(-\frac{4x}{y^5}\right) + (-p^2r^2)3xy^{1/2}$   
 $= 2(q-p) - 4pr^2s + \frac{8p^2rsx}{y^5} - 3p^2r^2xy^{1/2}$   
 $= -8y + \frac{(x+2y)x^3y^{1/2}(5x+2y)}{y^8}$

23.  $x^2 - xy + y^3 = 8$ , so let  $F(x, y) = x^2 - xy + y^3 - 8 = 0$ .  
Then  $\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{(2x-y)}{-x+3y^2} = \frac{y-2x}{3y^2-x}$ .

24.  $y^5 + 3x^2y^2 + 5x^4 = 12$ , so let  
 $F(x, y) = y^5 + 3x^2y^2 + 5x^4 - 12 = 0$ . Then  
 $\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{6xy^2 + 20x^3}{5y^4 + 6x^2y}$ .

25.  $x \cos y + y \cos x = 1$ , so let  
 $F(x, y) = x \cos y + y \cos x - 1 = 0$ . Then  
 $\frac{dy}{dx} = -\frac{\cos y - y \sin x}{-x \sin y + \cos x} = \frac{y \sin x - \cos y}{\cos x - x \sin y}$ .

26.  $2y^2 + \sqrt[3]{xy} = 3x^2 + 17$ , so let  
 $F(x, y) = 2y^2 + \sqrt[3]{xy} - 3x^2 - 17 = 0$ . Then  
 $\frac{dy}{dx} = -\frac{y / [3(xy)^{2/3}] - 6x}{4y + x / [3(xy)^{2/3}]} = \frac{18x - x^{-2/3}y^{1/3}}{12y + x^{1/3}y^{-2/3}}$ .

27. Let  $F(x, y, z) = xy + yz - xz = 0$ . Then  
 $\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{y-z}{y-x} = \frac{z-y}{y-x}$ ,  
 $\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{x+z}{y-x} = \frac{x+z}{x-y}$ .

28.  $x^2 + y^2 - z^2 = 2x(y + z)$ . Let

$$F(x, y, z) = x^2 + y^2 - z^2 - 2x(y + z) = 0. \text{ Then}$$

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{2x - 2y - 2z}{-2z - 2x} = \frac{x - y - z}{z + x},$$

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{2y - 2x}{-2z - 2x} = \frac{y - x}{z + x}$$

29.  $xy^2z^3 + x^3y^2z = x + y + z$ . Let

$$F(x, y, z) = xy^2z^3 + x^3y^2z - (x + y + z).$$

$$\text{Then } \frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{y^2z^3 + 3x^2y^2z - 1}{3xy^2z^2 + x^3y^2 - 1},$$

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{2xyz^3 + 2x^3yz - 1}{3xy^2z^2 + x^3y^2 - 1}.$$

30. Let  $F(x, y, z) = y^2ze^{x+y} - \sin(xyz) = 0$ . Then

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{y^2ze^{x+y} - yz \cos(xyz)}{y^2e^{x+y} - xy \cos(xyz)}$$

$$= \frac{z \cos(xyz) - yze^{x+y}}{ye^{x+y} - x \cos(xyz)}$$

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = \frac{xz \cos(xyz) - e^{x+y}(2yz + y^2z)}{y^2e^{x+y} - xy \cos(xyz)}$$

31.  $xy^2 + yz^2 + zx^2 = 3$ , so let

$$F(x, y, z) = xy^2 + yz^2 + zx^2 - 3 = 0.$$

$$\text{Then } \frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{y^2 + 2zx}{2yz + x^2} \text{ and}$$

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{2xy + z^2}{2yz + x^2}.$$

32. Let  $F(x, y, z) = xe^y + yz + ze^x = 0$ . Then

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{e^y + ze^x}{y + e^x}, \quad \frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{xe^y + z}{y + e^x}.$$

33.  $\ln(x + yz) = 1 + xy^2z^3$ , so let

$$F(x, y, z) = \ln(x + yz) - 1 - xy^2z^3 = 0. \text{ Then}$$

$$\begin{aligned} \frac{\partial z}{\partial x} &= -\frac{F_x}{F_z} = -\frac{1/(x + yz) - y^2z^3}{y/(x + yz) - 3xy^2z^2} \\ &= \frac{y^2z^3(x + yz) - 1}{y - 3xy^2z^2(x + yz)} \end{aligned}$$

$$\begin{aligned} \frac{\partial z}{\partial y} &= -\frac{F_y}{F_z} = -\frac{z/(x + yz) - 2xyz^3}{y/(x + yz) - 3xy^2z^2} \\ &= \frac{2xyz^3(x + yz) - z}{y - 3xy^2z^2(x + yz)} \end{aligned}$$

34.  $dr/dt = -1.2$ ,  $dh/dt = 3$ ,  $V = \pi r^2h$  and

$$dV/dt = 2\pi rh(dr/dt) + \pi r^2(dh/dt).$$

Thus when  $r = 80$  and  $h = 150$ ,

$$dV/dt = (-28,800)\pi + (19,200)\pi = -9600\pi \text{ cm}^3/\text{s}.$$